

AD-A061 028

NAVAL RESEARCH LAB WASHINGTON D C
ASTRON TYPE EQUILIBRIUM IN THE ABSENCE OF AN APPLIED MAGNETIC F--ETC(U)
NOV 78 C A KAPETANAKOS
NRL-MR-3886

F/G 20/9

UNCLASSIFIED

NL

OF
AD
A061028



END
DATE
FILMED
-79
DDC

ADA061028

DDC FILE COPY

P
nw

NRL Memorandum Report 3886


LEVEL

Astron Type Equilibrium in the Absence of an Applied Magnetic Field

C.A. KAPETANAKOS

*Experimental Plasma Physics Branch
Plasma Physics Division*

November 14, 1978



This research was supported by the Office of Naval Research
and the Department of Energy.



NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

78 11 06 122

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 3886	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ASTRON TYPE EQUILIBRIUM IN THE ABSENCE OF AN APPLIED MAGNETIC FIELD.		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem.
7. AUTHOR(s) C. A. Kapetanakos		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D.C. 20375		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem H02-28A
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) Ipp.		12. REPORT DATE November 1978
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. (14) NRL-MR-3886		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This research was supported by the Office of Naval Research and the Department of Energy.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Field reversed configuration Ion rings Astron device		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is shown that the Vlasov-Maxwell equations predict the existence of rotating layer equilibria (rigid rotor) in the absence of an external magnetic field, provided that the density of the layer exceeds a critical density.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251 950

LB

ASTRON TYPE EQUILIBRIUM IN THE ABSENCE OF AN APPLIED MAGNETIC FIELD

Using a cold fluid model, Yoshikawa¹ has shown the existence of equilibrium for a space charge neutralized, long rotating electron beam propagating in the absence of an externally applied magnetic field. In Yoshikawa's calculation the effect of the conducting wall surrounding the beam has been neglected. In recent experiments² at the Naval Research Laboratory, a rotating, overdense, space charge neutral relativistic electron beam has successfully propagated inside a conducting tube in the absence of an external magnetic field. In both the above two cases the azimuthal, self magnetic field (B_θ^s) of the propagating beam plays a significant role in the existence of equilibrium.

In this note, we make use of the steady state ($\frac{\partial}{\partial t} = 0$) Vlasov-Maxwell equations to show the existence of rotating layer (Astron type) equilibrium in the absence of an external magnetic field.

The azimuthally symmetric, z-independent equilibrium configuration is shown schematically in Fig. 1. It is assumed that the layer is space charge neutralized, its axial velocity $v_z = 0$ and the magnetic flux inside the conducting tube is conserved. The nonrelativistic, rotating layer is described by the distribution function

$$f_1^0(H - \omega P_\theta) = (\overline{m}\overline{n}/2\pi) \delta(H - \omega P_\theta - k_1), \quad (1)$$

where m, ω, \overline{n} and k_1 are constants. In addition, the total energy H

Note: Manuscript submitted October 4, 1978.

and the canonical angular momentum P_θ are constants of the motion. In the presence of an externally applied magnetic field, the equilibrium properties of proton (P)-layers that are described by the distribution function of Eq. (1) have been studied by Kapetanakis et al and their stability properties at a frequency near ω by Uhm and Davidson.⁴

Since $v_z = 0$, the argument of delta function in Eq. (3) may be expressed, in the nonrelativistic case, as

$$H - \omega P_\theta - k_1 = (m/2)[(v_\theta - r\omega)^2 + v_r^2] + U(r), \quad (2)$$

where

$$U(r) = -(qr/c)A_\theta^0(r)\omega - mr^2\omega^2/2 - k_1, \quad (3)$$

and A_θ^0 is the equilibrium magnetic vector potential that describes the axial, self magnetic field $B_z^s(r)$ of the rotating layer.

It is easy to show from the results of ref. 3, that in the absence of an external magnetic field B_0 , the density profile $n^0(r)$, the azimuthal current density $J_\theta^0(r)$, the mean azimuthal velocity of the rotating layer $V_\theta^0(r)$ the self magnetic field $B_z^s(r)$ and the inner a_1 and outer a_2 radii are given by

$$n^0(r) = \begin{cases} 0, & 0 \leq r < a_1, \\ \bar{n}, & a_1 \leq r \leq a_2, \\ 0, & a_2 < r < b, \end{cases} \quad (4)$$

$$J_\theta^0(r) = \begin{cases} 0, & 0 \leq r < a_1, \\ qnV_\theta^0(r), & a_1 \leq r \leq a_2, \\ 0, & a_2 < r \leq b, \end{cases} \quad (5)$$

$$v_{\theta}^0(r) = \langle v_{\theta}(r) \rangle = (1/\bar{n}) \int v_{\theta} f_1^0 d^2v = \omega r, \quad a_1 \leq r \leq a_2, \quad (6)$$

$$B_z^s(r) = \begin{cases} \frac{4\pi I_{\ell}}{c} \left[1 - \frac{(a_1^2 + a_2^2)}{2b^2} \right], & 0 \leq r < a_1, \\ \frac{4\pi}{c} I_{\ell} \left[\frac{(a_2^2 - r^2)}{(a_2^2 - a_1^2)} - \frac{(a_1^2 + a_2^2)}{2b^2} \right], & a_1 \leq r \leq a_2, \\ -\frac{4\pi I_{\ell}}{c} \left[\frac{(a_1^2 + a_2^2)}{2b^2} \right], & a_2 < r \leq b, \end{cases} \quad (7)$$

and

$$a_1^2 = - \frac{16k_1 \lambda^{*2}/(m\omega^2/b^2)}{1 \pm \left\{ 1 + 16(\lambda^*/b)^2 \left[1 + 4k_1/(m\omega^2 b^2) \right] \right\}^{\frac{1}{2}}} = - \frac{k_1 c^2}{\pi q I_{\ell} \omega}, \quad (8)$$

$$a_2^2 = b^2 + \frac{8\lambda^{*2} \left[1 + 2k_1/(m\omega^2 b^2) \right]}{1 \pm \left\{ 1 + 16(\lambda^*/b)^2 \left[1 + 4k_1/(m\omega^2 b^2) \right] \right\}^{\frac{1}{2}}} = b^2 + \frac{mc^2 \omega b^2 \left[1 + 2k_1/(m\omega^2 b^2) \right]}{2\pi q I_{\ell}}. \quad (9)$$

In these equations $I_{\ell} = (q\bar{n}\omega/2)(a_2^2 - a_1^2)$ is the azimuthal current per unit length, $\lambda^* = c/\omega_p$, $\omega_p^2 = 4\pi\bar{n}q^2/m$ and k_1 , I_{ℓ} and ω are negative for $q > 0$. Since $a_2^2 \leq b^2$ and $(a_2^2 - a_1^2)$ should be real, Eqs. (8) and (9) give

$$\frac{1}{2} \left[1 + \frac{b^2}{16\lambda^{*2}} \right] \leq \left(\frac{\omega_0}{\omega} \right)^2 \leq 1, \quad (10)$$

where $\omega_0^2 = 2|k_1|/mb^2$.

ACCESSION for	White Section	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	Buff Section	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
UNCLASSIFIED				
CLASSIFICATION				
DISTRIBUTION/AVAILABILITY CODES				
SPECIAL				
A				

According to Eq. (10), radially confined equilibria ($a_2 \leq b$) exist only for

$$b \geq 4\lambda^*, \quad (11)$$

i.e., for $\bar{n} = \bar{n}_{\min} \geq 4/\pi b^2 R_0$, where R_0 is the charged particle classical radius. For a rotating electron layer inside a conducting tube of radius $b = 4$ cm, Eq. (11) predicts that equilibrium exists only for densities above $3 \times 10^{11} \text{ cm}^{-3}$. For protons the minimum density is higher by the mass ratio m_p/m_e .

At the minimum density ($\bar{n}_{\min} = 4/\pi b^2 R_0$), $\omega^2 = \omega_0^2$ and the inner and outer radii of the layer are $a_1 = b/\sqrt{2}$ and $a_2 = b$. The allowed values of $(\omega/\omega_0)^2$ as a function of $b/4\lambda^*$ are shown in Fig. 2.

For $q > 0$, the function $U(r)$ of Eq. (3) is negative and has the property $U(a_1) = U(a_2) = 0$. The peak of the envelop function $U(r)$ occurs at a radius that can be determined from

$$\left. \frac{dU(r)}{dr} \right|_{r=\rho} = 0,$$

and is equal to

$$\rho^2 = (a_1^2 + a_2^2)/2. \quad (12)$$

Using Eqs. (7) and (11), it can be shown that

$$\omega = -\Omega(\rho) = -\frac{qB_z^s(\rho)}{mc}. \quad (13)$$

The magnetic field becomes equal to zero at the radius ρ_0 , that is easily determined from Eq. (7). If $v [= \frac{\pi \bar{n} q^2}{mc^2} (a_2^2 - a_1^2)]$ is the layer strength (Budker) parameter, Eq. (7) gives

$$\frac{(a_2^2 - \rho^2)}{(a_2^2 - a_1^2)} = \frac{(\nu + 1)}{2\nu} \quad (14)$$

Since $I_\ell = \nu mc^2 / 2\pi q$, the inner a_1 and outer a_2 radii can be written as

$$\frac{a_1^2}{b^2} = \left(\frac{\omega_0}{\omega} \right)^2 \frac{1}{\nu}, \quad (15)$$

and
$$\frac{a_2^2}{b^2} = \left(1 + \frac{1}{\nu} \right) - \frac{a_1^2}{b^2}, \quad (16)$$

substituting Eqs. (15) and (16) in Eq. (12), it is obtained

$$\frac{\rho^2}{b^2} = \frac{1}{2} \left(1 + \frac{1}{\nu} \right). \quad (17)$$

For $(\omega/\omega_0) = 1$, Eqs. (15) and (16) give $(a_2/b)^2 = 1$ and

$$(a_1/b)^2 = \frac{1}{2} \pm \frac{1}{2} \left(1 - \bar{n}_{\min}/\bar{n} \right)^{\frac{1}{2}}. \quad (18)$$

When $\bar{n} \gg \bar{n}_{\min}$, the only physically meaningful root of Eq. (18) is $(a_1/b)^2 = 0$, i.e., the layer becomes solid with a radius equal to the radius of the surrounding conducting tube.

In addition, for $\bar{n} \gg \bar{n}_{\min}$, Eq. (10) gives for the minimum allowed frequency of rotation ω

$$\left(\frac{\omega}{\omega_0} \right)^2 \cong 2\bar{n}_{\min}/\bar{n}. \quad (19)$$

Substituting Eq. (19) into Eqs. (15) and (16), we get $\frac{a_1}{b} = \frac{1}{2}$ and $\frac{a_2}{b} = \frac{\sqrt{3}}{2}$.

For $b \rightarrow \infty$, $\omega_0 \rightarrow 0$ and since $(a_2^2 - \rho^2)/(a_2^2 - a_1^2) = \frac{1}{2}$, the frequency of rotation ω remains $\neq 0$ for $I_\ell \neq 0$, as may be seen from Eqs. (7) and (13). Thus, according to Eq. (10), equilibrium does not exist.

In summary, I have shown that the Vlasov-Maxwell equations predict the existence of rigid-rotor type equilibria in the absence of an external magnetic field, provided that the particle density in the layer exceeds a minimum density given by $\bar{n}_{\min} = 4/\pi b^2/R_0^2$. For a proton layer confined inside a tube of $b = 10$ cm, $\bar{n}_{\min} = 8 \times 10^{13} \text{ cm}^{-3}$, which can be easily obtained with existing pulsed proton sources.⁵ However, for times longer than the ion-electron collision time the situation becomes more complicated because of the possibility of excitation of electron counter current.

References

1. S. Yoshikawa, Phys. Rev. Lett. 26, 295 (1971).
2. J. D. Sethian, K. A. Gerber, D. N. Spector and A. E. Robson, Phys. Rev. Lett. 41, 798 (1978).
3. C. A. Kapetanacos, J. Golden and K. R. Chu, Plasma Phys. 19, 387 (1977).
4. H. S. Uhm and R. C. Davidson, Stability Properties of a Field-Reversed Ion Layer in a Background Plasma, (unpublished).
5. J. Golden, C. A. Kapetanacos, S. J. Marsh and S. J. Stephanakis, Phys. Rev. Lett. 38, 130 (1977).




Fig. 1 — Equilibrium configurations of a long, rotating, space-charge neutral layer




Fig. 2 — Allowed region (cross hatched) of the normalized frequency ω/ω_0 as a function of $b/4\lambda^*$